

Effect of stellar activity on the high-precision transit light curve

Mahmoud Oshagh


Institute of Astrophysics and space science (IA)
Center of Astrophysics of Uni of Porto (CAUP)

Collaborators: N. Santos, D. Ehrenreich, N. Haghhighipour, G. Boué,
P. Figueira, A. Santerne, I. Boisse, X. Bonfils, M. Montalto


Outline

- Stellar activity and planetary transit
- Quantifying the spot occultation impact on our estimations
- Impact of stellar activity on transmission spectra
- Can stellar activity mimic the signature of blue sky in HD 189733b and GJ 3470b?
- Conclusions

Non-occulted stellar spots





$$\frac{\Delta f}{f} = \left(\frac{R_p}{R_*} \right)^2$$



CoRoT-2b (Czesla et al 2009)

Overestimation on the planet radius by 3–4% (Czesla et al 2009)

Occulted stellar spots



HAT-P-11b observed by Kepler

Occulted stellar spots

In the case that the spot anomaly is clearly identifiable




Some authors assigned a zero weight to the anomalous points of the light curve.



May not be the best approach, since the missing points in the transit light curve can cause offsets in the parameters estimation.






It is better to model them and then remove anomaly features from light curve.



HAT-P-11b observed by Kepler

SOAP-T

SOAP-T produces the expected light curve and the radial velocity signal of a system consisting of a rotating spotted star with a transiting planet.





Quantifying the impact

In case that the spot anomalies are not identifiable, we want to quantify their effect on our measurements.

Generated a large number of mock transit light curves with spot anomaly inside for different combination of planet radius and spot size


try to fit them with transit light curve without anomaly. Transit depth, duration and timing are free parameters



Transit timing measurements


Jupiter size planet transiting a solar-like star overlaps a spot with a filling factor of 1 %, produces the maximum value of TTV around **200 seconds**.

This amplitude of TTV can be induced by an Earth-mass planet in a mean-motion resonance with a transiting Jupiter-like planet on a 3 day orbit.





Transit timing measurements

Jupiter size planet transiting a solar-like star overlaps a spot with a filling factor of 1 %, produces the maximum value of TTV around **200 seconds**.




Transit timing measurements

TTV induced by stellar spot occultation may be used to distinguish between prograde and retrograde transiting planets (Mazeh et al 2014).




Planet radius estimations

A radius of Neptune size planet transiting a solar-like star and overlapping a spot with filling factor is 1 %, can be underestimated by 4%.




Transit duration estimation


A Jupiter size planet transiting a solar-like star and overlaps a spot with filling factor of 1% causes transit duration to be **4%, longer or shorter**, which affects the planetary inclination measurements.



Transmission spectroscopy




Multi-band photometry



HD 209458b from 290–1030 nm (Knutson et al. 2007)

- Molecular composition
- Thermal structure of the atmosphere

$$\frac{R_p}{R_*}(\lambda)$$





HD 209458b (Deming et al 2013)

Stellar activity impact on transmission spectra

Mostly the impact of non-occulted stellar spot are taking into account (e.g HD 189733b, GJ 3470b).

Occulted spots are corrected if their anomaly are clearly identified (e.g HD 189733b).




Quantifying the impact

Quantify the impact of occultation of stellar active regions, such as spot and plage, on the transmission spectroscopy measurements.


Modify SOAP-T to generate transit light curves at different wavelengths.

Generated a large number of mock transit light curves with spot anomaly inside for different combination of planet radius and spot/plage size.

Transit depth is the only free parameters, and compute the difference between fitted value and their actual value





Occultation with stellar spot



Occultation with stellar plage

Quantifying the impact


The anomalies inside the transit lead to a significant underestimation or overestimation of the planet-to-star radius ratio as a function of wavelength. At short wavelengths, the effect can reach up to a maximum difference of **10%** in the planet-to-star radius ratio.



HD 189733 b

HD 189733b is a hot Jupiter orbiting a K-dwarf. The brightness of HD 189733 ($V \simeq 7.7$) and the HD 189733b large planet to star radius ratio and also its short-orbit, has made it one of the first and also the most well studied planet in the planetary atmosphere investigations.


Sing et al. (2011) and Pont et al. (2013) both reported excess in the planet radius in the short wavelength (300–800nm) and the authors find a good agreement between this observation and the prediction of Rayleigh scattering in the planet atmosphere (blue sky).



Pont et al. (2013)

HD 189733 b

HD 189733 is an active star which shows photometric modulation up to $\approx 2\%$ during its 12 days stellar rotation period Boisse et al. (2009); Sing et al. (2011).





Pont et al. (2013)

HD 189733 b

We found out that the observed transmission spectrum of HD 189733b can be reproduced simply by considering the overlap of HD 189733b with a stellar plage with **filling factor of 1.96%** and a **temperature contrast of +100 K** with the stellar temperature.


Note: Determining the plage's temperature and its filling factor are strongly degenerate.



GJ 3470 b

GJ 3470b is a hot Uranus which orbits an active M-dwarf, GJ 3470,


An increment slope towards the blue side of spectrum ($\sim 360\text{nm}$), which was interpreted by Nascimbeni et al. (2013) as Rayleigh scattering processes in the planet's atmosphere (blue sky).



Nascimbeni et al. (2013)

GJ 3470 b



Fukui et al. (2013) reported photometric variability on GJ 3470 during 60 days observations around $\text{Photvar} \simeq 1\%$, which can be explained by GJ3470 being an active star which may harbor a spot with a filling factor of 1%.



Fukui et al. (2013)


GJ 3470 b

The plage which was needed to justify the observed excess planet radius has **filling factor around 2.56%** with a **temperature contrast of +100 K** with the stellar temperature. The size and relative brightness of the plage required by our scenario also generates a photometric variation around (1.3%) which is compatible with the observed one (Photovar $\simeq 1\%$) (Fukui et al. 2013).



Observational Strategy

- 1) Several observations of a transit in a given wavelength and use the variations of transit depth as a function of time
- 2) Simultaneous multi-band photometric observations for a quarter of the stellar rotation period before and after the transit.



Conclusions

In case of an active star we have to pay attention to the inside the transit light curves anomalies.

If we can identify spot occultation anomalies, we have to remove them by modeling them (e.g. using SOAP-T). If we could not identify them, we have to be careful about our planetary parameters estimation.

Prior to interpreting the transmission spectra and attempting to set constraints on the exoplanetary atmospheric models, it is critical to rule out the possible contamination due to stellar activity (both occulted and non-occulted stellar active region).

We showed that the transmission spectra of the active stars HD 189733b and GJ 3470b, and especially their excess of the planet radius in the bluer part of the spectra, can exactly be reproduced by assuming the occultation of the HD 189733b and GJ 3470b with the active region (plage) of their host star.

Thanks for your attention!

